

Publications 2002

Wild A, Jaeger M, Haak H, Mehdian SH.



Sacral insufficiency fracture, an unsuspected cause of low-back pain in elderly women.

Arch Orthop Trauma Surg. 2002 Feb;122(1):58-60.

Agorastides ID, Lam KS, Freeman BJ, Mulholland RC.



The Adams classification for cadaveric discograms: inter- and intra-observer error in the clinical setting.

Eur Spine J. 2002 Feb;11(1):76-9.

Debnath UK, De la Harpe D, Webb JK.



A complex thoracic fracture extending from fourth thoracic to twelfth thoracic vertebrae without neurological deficit--a case report.

Injury. 2002 Mar;33(2):188-90. No abstract available.

Freeman BJ, Oullet JA, Webb JK.



Excision of hemivertebrae in the management of congenital scoliosis.

J Bone Joint Surg Br. 2002 Mar;84(2):305; author reply 306. No abstract available.

Schaan M, Jaksche H, Boszczyk B.



Predictors of outcome in head injury: proposal of a new scaling system.

J Trauma. 2002 Apr;52(4):667-74.

Boszczyk AA, Boszczyk BM, Putz RV.



Prenatal rotation of the lumbar spine and its relevance for the development of the zygapophyseal joints.

Spine. 2002 May 15;27(10):1094-101.

Pratt RK, Burwell RG, Cole AA, Webb JK.



Patient and parental perception of adolescent idiopathic scoliosis before and after surgery in comparison with surface and radiographic measurements. *Spine*. 2002 Jul 15;27(14):1543-50; discussion 1551-2.

Pelosi L, Lamb J, Grevitt M, Mehdian SM, Webb JK, Blumhardt LD.



Combined monitoring of motor and somatosensory evoked potentials in orthopaedic spinal surgery. *Clin Neurophysiol*. 2002 Jul;113(7):1082-91.

Pratt RK, Webb JK, Burwell RG, Cole AA.



Re: Koch et al. Adolescents undergoing surgery for idiopathic scoliosis: how physical and psychological characteristics relate to patient satisfaction with cosmetic result. (*Spine* 2001;26:2119-24). *Spine*. 2002 Jul 15;27(14):1594-5; author reply 1594-5. No abstract available.

Sengupta DK, Mehdian SH, McConnell JR, Eisenstein SM, Webb JK.



Pelvic or lumbar fixation for the surgical management of scoliosis in duchenne muscular dystrophy. *Spine*. 2002 Sep 15;27(18):2072-9.

Wild A, Jäger M, Koch H, Webb JK.



Treatment of congenital scoliosis in an 8-month-old child. *Arch Orthop Trauma Surg*. 2002 Sep;122(7):418-20. Epub 2002 Jul 3.

Boszczyk B, Bierschneider M, Potulski M, Robert B, Vastmans J, Jaksche H.



[Extended kyphoplasty indications for stabilization of osteoporotic vertebral compression fractures] *Unfallchirurg*. 2002 Oct;105(10):952-7. German.

Mulholland RC, Sengupta DK.



Rationale, principles and experimental evaluation of the concept of soft stabilization.

Eur Spine J. 2002 Oct;11 Suppl 2:S198-205. Epub 2002 Jun 4. Review.

Sengupta DK, Mehdian SM, Mulholland RC, Webb JK, Ohnmeiss DD.



Biomechanical evaluation of immediate stability with rectangular versus cylindrical interbody cages in stabilization of the lumbar spine.
BMC Musculoskelet Disord. 2002 Oct 3;3:23.

Benjamin M, Kumai T, Milz S, Boszczyk BM, Boszczyk AA, Ralphs JR.



The skeletal attachment of tendons--tendon "enthyses".
Comp Biochem Physiol A Mol Integr Physiol. 2002 Dec;133(4):931-45.
Review

Webb JK.



Comment to "Clinical outcome and radiographic results after operative treatment of Scheuermann's disease", by R. W. Poolman et al.
Eur Spine J. 2002 Dec;11(6):570. No abstract available

Burwell RG, Aujla KK, Cole AA, Kirby AS, Pratt KK, Webb JK, Moulton A.



Anterior universal spine system for adolescent idiopathic scoliosis: a follow-up study using scoliometer, real-time ultrasound and radiographs.
Stud Health Technol Inform. 2002;91:473-6.

Burwell RG, Aujla RK, Cole AA, Kirby AS, Pratt RK, Webb JK, Moulton A.



Preliminary study of a new real-time ultrasound method for measuring spinal and rib rotation in preoperative patients with adolescent idiopathic scoliosis.
Stud Health Technol Inform. 2002;91:262-6.

Burwell RG, Aujla RK, Cole AA, Kirby AS, Pratt RK, Webb JK, Moulton A.



Spine-rib rotation differences at the apex in preoperative patients with adolescent idiopathic scoliosis: evaluation of a three-level ultrasound method.
Stud Health Technol Inform. 2002;91:246-50.

Burwell RG, Aujla RK, Cole AA, Kirby AS, Pratt RK, Webb JK, Moulton A.



Back shape assessment in each of three positions in preoperative patients with adolescent idiopathic scoliosis: evaluation of a 10-level Scoliometer method interpolated to 18-levels.

Stud Health Technol Inform. 2002;91:119-22.

Burwell RG, Pratt RK, Webb JK.



Etiologic theories of idiopathic scoliosis: the apical spinal deformity--relevance to surgical practice.

Stud Health Technol Inform. 2002;91:32-6

Burwell RG, Dangerfield PH.



Etiologic theories of idiopathic scoliosis: neurodevelopmental concepts to be evaluated.

Stud Health Technol Inform. 2002;91:15-9.

Burwell RG, Dangerfield PH.



The NOTOM hypothesis for idiopathic scoliosis: is it nullified by the delayed puberty of female rhythmic gymnasts and ballet dancers with scoliosis?

Stud Health Technol Inform. 2002;91:12-4.

Burwell RG.



Biology is the future of scoliosis treatment.

Stud Health Technol Inform. 2002;88:309-15. Review. No abstract available.

Burwell RG, Aujla RK, Kirby AS, Moulton A, Webb JK.



The early detection of adolescent idiopathic scoliosis in three positions using the scoliometer and real-time ultrasound: should the prone position also be used?

Stud Health Technol Inform. 2002;88:74-80.